

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re PATENT APPLICATION OF

Confirmation No. 7969

Darrell R. ANDERSON et al.

Group Art Unit: 1644

Application Serial No. 10/073,138

Examiner: Phillip Gambel

Filed: February 13, 2002

Title: IDENTIFICATION OF UNIQUE BINDING INTERACTIONS BETWEEN CERTAIN

ANTIBODIES AND THE HUMAN B7.1 AND B7.2 CO-STIMULATORY ANTIGENS

SUPPLEMENTAL INFORMATION DISCLOSURE STATEMENT

Commissioner for Patents P. O. Box 1450 Alexandria, VA 22313-1450

Sir:

Pursuant to 37 C.F.R. §.1.56, the attention of the Patent and Trademark Office is hereby directed to the references listed on the attached Form PTO-1449. Unless otherwise indicated herein, one copy of each reference is attached. It is respectfully requested that the listed references be expressly considered during the prosecution of this application, and that the references be made of record therein and appear among the "References Cited" on any patent to issue there from.

This Information Disclosure Statement is being filed before the mailing date of the first official action on the merits in the present application. No certification or fee is required, pursuant to 37 C.F.R. §.1.97(b)(3),

Respectfully Submitted,

Thomas A. Cawley, Jr., Ph.D.

-A Cowy /

Registration Number 40,944

Date: October 19, 2004

PILLSBURY WINTHROP LLP

Telephone: (703) 905-2000 Facsimile: (703) 905-2500

P.O. Box 10500 McLean, VA 22102

TAC/CR/sm

FORM PT	O-1449		dified) .	OCT 1 9 2004	FIGE LE	-	Atty. Ref. No 037003-0280			ent R 95-30	ef. -02330	CP3			
INFORMA BY APPL		DISC	LOSURE ST	TEMENT TRADE			Applicant:	ANDERSON	et al.						
							Appln. No.:	10/073,138							
							Filing Date:	February 13,	2002						
Date: Octo	ber 15	, 200	4	Page 1 of 2			Examiner:	Gambel, P.	Group Art	Unit:	164	14			
U.S. PATI	ENT DO	CUN	MENTS												
Examiner's	Do		cument	Date	ite Name			Class	Sub	Sub		Filing Date			
Initials*	Nu		nber	MM/YYYY	(Family Na	me of	First Inventor	First Inventor)			ss	(if appro	priate)		
	AR		5,304,635	04/1994	Imam	lmam				↓	 				
	BR	-					-,								
<u> </u>	CR					*									
FOREIGN	PATE	NT D	OCUMENTS			•	,	. *	English Abstract			lation ily Ava	ilable		
		Doc	ument		Country	Inve	ntor Name								
		Num	ber	MM/YYYY		1				Г	<u> </u>				
	1.					 	 		Enclosed	No	Enclo	se	No		
	DR					-			<u> </u>	-	-		┼		
	ER	<u> </u>				 			-	╁	┼		┼		
OTUED (FR		L' 1 A - 41	TH. D.	is dis al Mana	<u> </u>	- Dartin - A D			1	 		┼		
OTHER (I	nciudin GR	g in t					e, Pertinent P mutation on t	rages, etc.) the activating a	_l nd immunos	uppr	<u> </u>	prope	rties		
			of a "Human	ized" OKT3 r	nonoclonal a	ntiboo	dy," J. Immund	ol., 1992, 148:3	461-3468.						
	HR		Azuma M., et al., "CD28 Interaction with B7 Costimulates Primary Allogeneic Proliferative Responses and Cytotoxicity Mediated by Small Resting T Lymphocytes," J. Exp. Med., 1992, 175:353-360. Azuma M., et al., "Functional Expression of B7/BB1 on Activated T Lymphocytes," J. Exp. Med., 1993, 177:845-850.												
IR															
	JR		Azuma M.D., et al., "B70 Antigen is a Second Ligand for CTLA-4 and CD28," Nature, 1993, 366:76-79.												
KR			Boussiotis V.A., et al., "Activated human B lymphocytes express three CTLA-4 counter-receptors that costimulate T-Cell activation," Proc. Natl. Acad. Sci., USA, 1993, 90:11059-11063.												
	LR		Cohen J., "Mounting a targeted strike on unwanted immune responses," (news; comment), Science, 1992, 257:751.												
	MR		De Boer M.,	•	ctional Characterization of a Novel Anti-B7 Monoclonal Antibody," 071-3075.							Journa	al of		
	NR	is the CD86 differentiation antigen," Blood, 1994, 84, 1402-1407.													
	OR									nphoc	ytes				
	PR									on," Ti	ne				
	QR						_	nmune Toleran	ce," Immun	ology	Today	, 1993	3,		
	RR		Gimmi C.D.,				nergy is Induc 3, 90:6586-65	ced by Antigen I	Presentation	n in th	ne Abs	ence o	of B7		
	SR	·		., et al., "CTL				poptosis of hun	nan T cells.	' Prod	c. Natl.	Acad	. Sci.		
	TR		Grumet F.C., et al., "Soluble form of an HLA-B7 Class I Antigen Specifically Suppresses Humoral Alloimmunization." Human Immunology, 1994, 40:228-234.												
	UR		Harding F.A.		28 Mediated	Signa	alling Co-stimu	ulates Murine T	Cells and F	reve	nts Ind	uction	of		
	VR		Hart D.N.J.,		3-1 is a Leuc	ocyte	Differentiation	n Antigen on Hu	ıman Dendi	itic C	ells In	duced	by		
	WR		Ionescu-Tirg	oviste, et al,	"Correlations	s betw		ntibodies and th	ne HLA syst	em in	a grou	up of T	уре І		

	Janeway C.A., Jr. and K. Bottomly, "Signals and Signs for Lymphocyte Responses," Cell, 1994, 76.2 285.							
YR	Jenkins M.K., "The Role of Cell Division in the Induction of Clonal Anergy." Immunology Today, 1992, 13:69.							
ZR	June C.H., et al., "The B7 and CD28 receptor families," Immunol Today, 1994, 15:321-331.							
AAR	Krummel M., et al., "CD28 and CTLA-4 have opposing effects on the response of T cells to stimulation," J. Exp. Med. 1995, 182:459-466.							
SIPE BBR	LaSalle J.M., et al., "Early signaling defects in human T cells anergized by T cell presentation of autoantigen," J. Exp. Med., 1992, 176:177-186.							
nct 1 9 mix CB	Lenschow D.J., et al., "Long-Term Survival of Xenogeneic Pancreatic Islet Grafts Induced by CTLA-4Ig," Science, 1992, 257:789-795.							
DE THANK EER	Lenschow D.J., et al., "Expression and Functional Significance of an Additional Ligand for CTLA-4," F Natl. Acad. Sci., USA,,1993, 90:11054-11058.							
EER	Lin H., et al., "Long-term Acceptance of Major Histocompatibility Complex Mismatched Cardiac Allograf Induced by CTLA-4-Ig Plus Donor Specific Transfusion," J. Exp. Med., 1993, 178:1801.							
FFR	Linsley, P.S., et al., "CTLA-4 is a Second Receptor for the B Cell Activation Antigen B7," J. Exp. Med., 1991, 174:561.							
GGR	Linsley P.S., et. al., "T-Cell Antigen CD28 Mediates Adhesion with B Cells by Interacting with Activa Antigen B7/BB-1." Proc. Natl. Acad. Sci., 1990, 87:5031-5035.							
HHR	Linsley, et al, "CD28 Engagement by B7/BB-1 Induces Transient Down-Regulation of CD28 Synthes Prolonged Unresponsiveness to CD28 Signaling," The Journal of Immunology, 1993, 150:3161-3169							
IIR	Linsley, et al., "Binding of the B cell activation antigen B7 to CD28 costimulates T cell proliferation and interleukin 2 mRNA accumulation," J. Exp. Med., 1991, 173:721-730.							
JJR	Linsley P.S., et. al., "Coexpression and Functional Cooperation of CTLA-4 and CD28 on Activated T Lymphocytes." J. Exp. Med., 1992, 176:1595-1604.							
KKR	Morton P.A., et al., "Differential effects of CTLA-4 substitutions on the binding of human CD80 (B7-1) and CD86 (B7-2)," J. Immunol., 1996,156:1047-1054.							
LLR	Nestle F.O., et al, "Characterization of dermal dendritic cells in psoriasis," J. Clin. Invest., 1994, 94: 202-209.							
MMR	Schwartz R.H., "Co-stimulation of T lymphocytes: The role of CD28, CTLA-4, and B7/BB1 in interleukin-production and immunotherapy," Cell, 1992, 71:1065-1068.							
NNR	Schwartz R.H., "A cell culture model for T lymphocyte clonal anergy," Science, 1990, 248:1349-1356.							
OOR	Selvakumar, A., et al., "Genomic organization and chromosomal location of the human gene encoding th B-lymphocyte activation antigen B7," lmmunogenetics , 1992, 36:175-181							
PPR	Tan P., et al., "Induction of Alloantigen-specific Hyporesponsiveness in Human T Lymphocytes by Block Interaction of CD28 with Its Natural Ligand B7/BB1," J. Exp. Med., 1993, 177:165-173.							
QQR	Tivol E.A., et al., "Loss of CTLA-4 leads to massive lymphoproli-feration and fatal multiorgan tissue destruction, revealing a critical negative regulatory role of CTLA-4," Immunity, 1995, 3:541-547.							
RRR	Turka L.A., et al., "T-cell activation by the CD28 ligand B7 is required for cardiac allograft rejection in vivo," Proc. Natl. Acad. Sci., USA, 1992, 89:11102-11105.							
SSR	Toubert A., et al., "Epitope mapping of an HLA-B27 monoclonal antibody that also reacts with a 35-kD bacterial out-membrane protein," Clin. Exp. Immunol., 1990, 82(1), 16-20.							
TTR	Toubert A., et al., "Epitope mapping of HLA-B27 and HLA-B7 antigens by using intradomain recombinants, J. Immunol., 1988, 141(7), 2503-9.							
UUR	Valle et al., "mAb 104, a new monoclonal antibody, recognizes the B7 antigen that is expressed on activated B cells and HTLV-1-transformed T cells," lmmunol. , 1990, 69(4), 531-535.							
VVR	Vandenberghe P., et al "Antibody and B7/BB1-mediated ligation of the CD28 receptor induces tyrosine phosphorylation in human T cells," J. Exp. Med., 1992, 175:951-960.							
WWR	van der Merwe, et al., "CD80 (B7-1) binds both CD28 and CTLA-4 with a low affinity and very fast kinetics," J. Exp. Med., 1997,185: 393-403.							
XXR	Weyl D., et al., "Epitope mapping of human monoclonal antibodies to HLA-B27 by using natural and mutated antigenic variants," Hum. Immunol., 1991, 31(4), 271-276.							
YYR	Zavazava N., et al, "Inhibition of anti-HLA-B7 alloreactive CTL by affinity-purified soluble HLA," Transplantation, 1991, 51(4), 838-42.							

*EXAMINER: Initial if citation considered, whether or not citation is in conformance with MPEP § 609. Draw line through citation if not in conformance and not considered. Include copy of this form with next communication to Applicant.

PAT-1449 12/98